



Draft Genome Sequence of *Sphingobacterium* sp. CZ-UAM, Isolated from a Methanotrophic Consortium

José Luis Steffani-Vallejo,^a Cristal Zuñiga,^{b,c} Pablo Cruz-Morales,^a Luis Lozano,^d Marcia Morales,^b Cuauhtemoc Licona-Cassani,^{a,e} Sergio Revah,^b José Utrilla^d

StrainBiotech, Parque de Innovación AGROBIOTEC, Irapuato, Guanajuato, Mexico^a; Departamento de Procesos y Tecnología, Universidad Autónoma Metropolitana-Cuajimalpa, Ciudad de México, Mexico^b; Department of Pediatrics, University of California, San Diego, La Jolla, California, USA^c; Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico^d; Centro de Biotecnología-FEMSA, Tecnológico de Monterrey, Monterrey, Nuevo León, Mexico^e

ABSTRACT *Sphingobacterium* sp. CZ-UAM was isolated from a methanotrophic consortium in mineral medium using methane as the only carbon source. A draft genome of 5.84 Mb with a 40.77% G+C content is reported here. This genome sequence will allow the investigation of potential methanotrophy in this isolated strain.

Methane (CH₄) is produced by natural and human-related activities; it is one of the main greenhouse gases, with a 25-fold greater global warming potential than CO₂ (1). CH₄-consuming bacteria are known as methanotrophs, and they can grow using CH₄ as their only carbon source. Biological treatments are the best option to eliminate CH₄ as an atmospheric contaminant when its percentage is below the explosive limit (5% in air) (2, 3). There are many sources of cheap and diluted CH₄ that can be used as feedstock (4). Therefore, it is environmentally urgent to study new methanotrophs with higher CH₄ uptake rates and to explore their metabolic potential to convert CH₄ into added-value products (3, 5, 6). Here, we report the genome sequence of the novel potential methanotroph *Sphingobacterium* sp. CZ-UAM, which was isolated from a methanotrophic consortium from a wastewater treatment plant (7). The isolated strain is able to grow on mineral medium with a 5% CH₄ atmosphere as its sole carbon source.

Genomic DNA from *Sphingobacterium* sp. CZ-UAM was extracted by an in-house phenol-chloroform protocol, RNase treated, and gel verified for quality. Then, a paired-end (average size, 527 bp) genomic DNA library was generated using commercial kits according to the manufacturers' instructions (TruSeq Nano, Illumina, San Diego, CA, USA). The DNA library quality was verified using a 2100 BioAnalyzer (Agilent Genomics, Santa Clara, CA, USA). The genome sequencing was performed using an Illumina HiSeq 2500 system through the standard rapid sequencing protocol for cluster generation and sequencing by synthesis, which resulted in a total of 3,891,197 reads for a depth coverage of 300×. Genome assembly was performed using SPAdes version 3.9.0 (8) and IDBA_UD version 1.1.1 (9). Then, SSPACE version 3.0 (10) and Metassembler version 1.5 (11) were used for merging and scaffolding the obtained contigs. After genome assembly and scaffolding, 12 contigs were obtained. Phylogenetic analysis of the 16S rRNA was performed by extracting the previously annotated 16S ribosomal gene sequence, as well as the annotated *Sphingobacterium* sequences in the NCBI database. All the sequences were aligned using Muscle software (12) and edited using Gblocks (13) prior to phylogenetic tree construction using Bayesian inference. The analyses indicate that the isolated strain is a member of the *Sphingobacterium* genus. The analysis of the full 16S rRNA gene sequence with the EZBioCloud database indicates

Received 27 June 2017 Accepted 3 July 2017 Published 17 August 2017

Citation Steffani-Vallejo JL, Zuñiga C, Cruz-Morales P, Lozano L, Morales M, Licona-Cassani C, Revah S, Utrilla J. 2017. Draft genome sequence of *Sphingobacterium* sp. CZ-UAM, isolated from a methanotrophic consortium. Genome Announc 5:e00792-17. <https://doi.org/10.1128/genomeA.00792-17>.

Copyright © 2017 Steffani-Vallejo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to José Utrilla, utrilla@ccg.unam.mx.

higher identity with *S. pakistanense* (97.1%) and *S. canadense* (96.8%), *S. ginsenosidimutans* (96.46%), and *S. detergens* (95.39%) (14). Of the few available *Sphingobacterium* sequencing projects, none of them have been reported as having the potential to be a methanotroph. The availability of the *Sphingobacterium* sp. CZ-UAM genome will contribute to the study of the *Sphingobacterium* genus and will provide information on the molecular basis of this strain's ability to consume CH₄ as a sole carbon source.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession no. [MTCM0000000](https://doi.org/10.1093/cmb.2012.0021). The version described in this paper is the first version, MTCM0100000 (BioSample SAMN06215933).

ACKNOWLEDGMENTS

We thank Alfredo Hernández and Víctor Del Moral for their computational support. Partial support from both UNAM-PAPITT-DGAPA (grant IA200716) and CONACyT (grant PN2015-01-241) is acknowledged.

REFERENCES

- Goodland R, Anhang J. 2009. Livestock and climate change: what if the key actors in climate change are...cows, pigs, and chickens? Worldwatch Institute, Washington, DC.
- Rocha-Rios J, Muñoz R, Revah S. 2010. Effect of silicone oil fraction and stirring rate on methane degradation in a stirred tank reactor. *J Chem Technol Biotechnol* 85:314–319. <https://doi.org/10.1002/jctb.2339>.
- Zuñiga C, Morales M, Le Borgne S, Revah S. 2011. Production of poly-β-hydroxybutyrate (PHB) by *Methylobacterium organophilum* isolated from a methanotrophic consortium in a two-phase partition bioreactor. *J Hazard Mater* 190:876–882. <https://doi.org/10.1016/j.jhazmat.2011.04.011>.
- Strong PJ, Xie S, Clarke WP. 2015. Methane as a resource: can the methanotrophs add value? *Environ Sci Technol* 49:4001–4018. <https://doi.org/10.1021/es504242n>.
- Wendlandt K-D, Stottmeister U, Helm J, Soltmann B, Jechorek M, Beck M. 2010. The potential of methane-oxidizing bacteria for applications in environmental biotechnology. *Eng Life Sci* 10:87–102. <https://doi.org/10.1002/elsc.200900093>.
- Zuñiga C, Morales M, Revah S. 2013. Polyhydroxyalkanoates accumulation by *Methylobacterium organophilum* CZ-2 during methane degradation using citrate or propionate as cosubstrates. *Bioresour Technol* 129:686–689. <https://doi.org/10.1016/j.biortech.2012.11.120>.
- Rocha-Rios J, Bordel S, Hernández S, Revah S. 2009. Methane degradation in two-phase partition bioreactors. *Chem Eng J* 152:289–292. <https://doi.org/10.1016/j.cej.2009.04.028>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Peng Y, Leung HCM, Yiu SM, Chin FYL. 2012. IDBA-UD: a *de novo* assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* 28:1420–1428. <https://doi.org/10.1093/bioinformatics/bts174>.
- Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27:578–579. <https://doi.org/10.1093/bioinformatics/btq683>.
- Wences AH, Schatz MC. 2015. Metassembler: merging and optimizing *de novo* genome assemblies. *Genome Biol* 16:207. <https://doi.org/10.1186/s13059-015-0764-4>.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797. <https://doi.org/10.1093/nar/gkh340>.
- Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 17:540–552. <https://doi.org/10.1093/oxfordjournals.molbev.a026334>.
- Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J. 2017. Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* 67:1613–1617. <https://doi.org/10.1099/ijsem.0.001755>.